

*Annual Review of Genetics*

# Regulation by 3'-Untranslated Regions

Christine Mayr

Department of Cancer Biology and Genetics, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA; email: mayrc@mskcc.org

Annu. Rev. Genet. 2017. 51:171–94

First published as a Review in Advance on August 30, 2017

The *Annual Review of Genetics* is online at [genet.annualreviews.org](http://genet.annualreviews.org)

<https://doi.org/10.1146/annurev-genet-120116-024704>

Copyright © 2017 by Annual Reviews.  
All rights reserved

## Keywords

noncoding RNA, regulatory RNA, alternative 3'-UTRs, alternative polyadenylation, cotranslational protein complex formation, cellular organization, mRNA localization, RNA-binding proteins, cooperativity, accessibility of regulatory elements

## Abstract

3'-untranslated regions (3'-UTRs) are the noncoding parts of mRNAs. Compared to yeast, in humans, median 3'-UTR length has expanded approximately tenfold alongside an increased generation of alternative 3'-UTR isoforms. In contrast, the number of coding genes, as well as coding region length, has remained similar. This suggests an important role for 3'-UTRs in the biology of higher organisms. 3'-UTRs are best known to regulate diverse fates of mRNAs, including degradation, translation, and localization, but they can also function like long noncoding or small RNAs, as has been shown for whole 3'-UTRs as well as for cleaved fragments. Furthermore, 3'-UTRs determine the fate of proteins through the regulation of protein–protein interactions. They facilitate cotranslational protein complex formation, which establishes a role for 3'-UTRs as evolved eukaryotic operons. Whereas bacterial operons promote the interaction of two subunits, 3'-UTRs enable the formation of protein complexes with diverse compositions. All of these 3'-UTR functions are accomplished by effector proteins that are recruited by RNA-binding proteins that bind to 3'-UTR *cis*-elements. In summary, 3'-UTRs seem to be major players in gene regulation that enable local functions, compartmentalization, and cooperativity, which makes them important tools for the regulation of phenotypic diversity of higher organisms.



### ANNUAL REVIEWS **Further**

Click [here](#) to view this article's online features:

- Download figures as PPT slides
- Navigate linked references
- Download citations
- Explore related articles
- Search keywords

## INTRODUCTION


Regulation of gene function through 3'-untranslated regions (3'-UTRs) is a relatively new field as only recent sequencing technology has provided us with the full landscape of 3'-UTRs across species and cell types. Before sequencing technology was available, detailed functional and mechanistic studies were performed only on a few model 3'-UTRs. Although these model 3'-UTRs have contributed substantially to our understanding of 3'-UTR biology, the conclusions drawn about their regulatory functions have focused primarily on the role of 3'-UTRs in gene regulation. 3'-UTRs determine protein levels through regulation of mRNA stability and translation mediated largely by AU-rich elements and miRNAs (6, 7, 21). 3'-UTRs also enable local translation through the regulation of mRNA localization (82, 98). In this review, some of the general principles that were learned from these model 3'-UTRs will be discussed, but I mostly focus on newer findings that may change our view about the functions of 3'-UTRs. This includes the discovery that 3'-UTR length can be regulated by alternative cleavage and polyadenylation (APA) (81, 87, 106), as well as the observation that 3'-UTRs may be cleaved off and act like long noncoding RNAs living their own life independent of the coding region (17, 69, 90). Lastly, 3'-UTRs were found to be able to mediate protein-protein interactions (PPIs), a discovery which has widespread consequences for protein complex formation, protein localization, and protein functions (9, 19, 31, 45).

3'-UTRs regulate gene expression through the binding of RNA-binding proteins (RBPs) (5, 15, 51, 93). RBPs bind to 3'-UTR *cis*-elements and mediate 3'-UTR functions through the recruitment of effector proteins. As RBPs interact with diverse effector proteins, each 3'-UTR regulatory element has the potential to carry out several different functions, depending on the cell type or cellular state. The cellular state also determines the RBPs that are able to access 3'-UTRs at a given moment. As such, 3'-UTR functions can be assessed only in the context of their bound RBPs. The composition of RBPs bound to a 3'-UTR at a given moment is dynamic and can change depending on the local environment, e.g., through addition of posttranslational modifications (PTMs), local expression of other RBPs, and interactions with membranes and cytoskeletal filaments (59, 60, 118). RBP binding is also influenced by secondary and tertiary RNA structure formation that regulates the accessibility of 3'-UTRs (2, 42, 68, 122). Importantly, RBPs cooperate with other bound RBPs to enable functional specificity *in vivo*. Answers to open questions will ultimately stimulate new avenues of 3'-UTR research in coming years.

In 1958, Francis Crick wrote, "Watson said to me, a few years ago, 'The most significant thing about the nucleic acids is that we don't know what they do.' By contrast the most significant thing about proteins is that they can do almost anything" (24, p. 138). In the same article, Crick published his hypothesis of the central dogma, in which he stated that the "transfer of information from nucleic acid to nucleic acid, or from nucleic acid to protein may be possible, but transfer from protein to protein, or from protein to nucleic acid is impossible" (24, p. 153). The simplicity of the idea was widely embraced and shaped the protein-centric view of cell biology for the next 50 years. However, in addition to the coding region—which is translated into protein—mRNAs also consist of untranslated regions at the 5' and 3' ends, which contain important regulatory elements (103). The novel finding that the nucleic acids located within 3'-UTRs can mediate PPIs, and thus can determine protein functions without altering the amino acid sequence (9), has the potential to lead to a less protein-centric view of cell biology.

## EXPANSION OF 3'-UTRs DURING EVOLUTION OF MORE COMPLEX ORGANISMS

3'-UTRs are ancient components of mRNAs and are found in all three domains of life—bacteria, archaea, and eukaryotes. Despite the compact genomes found in unicellular organisms, sequencing



**Figure 1**

Characteristics of 3'-UTR isoforms. (a) Correlation between genome size and median 3'-UTR length in uni- and multicellular organisms. Genome size was obtained from Reference 97. 3'-UTR length was obtained from References 26–28, 58, 81, 107, 130, and 139. (b) Sequence conservation shown as PhastCons scores for genes that generate three 3'-UTRs ( $n = 1,939$ ) for the regions between the stop codon and the first pA site (short 3'-UTR), the first to the second pA site (middle 3'-UTR), and the second to the last pA site (distal 3'-UTR). (c) Alternative 3'-UTRs are expressed in a cell type-specific manner. Shown are 3'-seq data for skeletal muscle (*top*) and naive B cells (*bottom*) obtained from Reference 81. The gene model (*gray*) is drawn to scale and depicts the last exon. Abbreviations: 3'-UTR, 3'-untranslated region; Mbp, millions of base pairs; nt, nucleotide; pA, polyadenylation.

technology has revealed that their mRNAs contain 3'-UTRs (26, 27, 107, 139). In unicellular genomes, medium 3'-UTR length correlates with genome size (**Figure 1a**) and is approximately 60 nucleotides (nt) in the bacterial and archaeal species sequenced so far and approximately 150 nt in yeast. Characteristic of the evolution of more complex multicellular organisms is a substantial expansion in genome size that also strongly correlates with increased 3'-UTR length (28, 58, 81, 130) (**Figure 1a**). For example, median 3'-UTR length increased from approximately 140 nt in worms to approximately 1,200 nt in humans (86). Intriguingly, the increase in 3'-UTR length correlates with organismal complexity when measured as the number of cell types found in an organism (20).

Notably, although they are less conserved than coding regions, 3'-UTRs still show a high degree of sequence conservation across species (117, 143), which is exemplified by the distribution of highly conserved elements. The percentage of base pairs in vertebrate genomes that are highly conserved is 0.14%; however, these elements are enriched 22-fold in coding regions and 11-fold in 3'-UTRs (117). Furthermore, 3'-UTRs often contain many islands of sequence conservation that are similar in conservation to coding regions, and these islands often contain binding sites for miRNAs or RBPs (34, 143). Taken together, the increase in 3'-UTR length during the

development of higher organisms and the enrichment of conserved sequence elements within 3'-UTRs suggests a critical role for 3'-UTRs in the regulation of biological complexity (86).

### ALTERNATIVE CLEAVAGE AND POLYADENYLATION REGULATES THE PRESENCE AND ACCESSIBILITY OF REGULATORY ELEMENTS IN 3'-UTRs

Further evidence for the importance of 3'-UTRs is the fact that the presence of functional elements is regulated through the process of APA. In archaea, 30% of genes generate alternative 3'-UTRs with a median length difference of 54 nt (26). In humans, the median 3'-UTR length of genes that produce alternative 3'-UTR isoforms is 2,462 nt, and the majority of genes (51–79%) express alternative 3'-UTRs (28, 81; I. Singh & C. Mayr, unpublished data). Thus, the number of genes that generate alternative 3'-UTRs has expanded alongside the increase in 3'-UTR length during the evolution of multicellular organisms.

Intuitively, one would think that longer 3'-UTRs mediate more regulatory functions as they contain more regulatory elements. However, this is only partially the case, as it was found that especially long 3'-UTRs are less responsive than expected to miRNA regulation (2, 68). One reason for this relationship seems to be a decreased accessibility of regulatory elements because of occlusive structures in long 3'-UTRs. As was shown several years ago, miRNA-binding sites located in the middle of 3'-UTRs mediate less repression than do sites at the ends of 3'-UTRs (42), which is also true for alternative 3'-UTRs (50). Better accessibility is supported by cross-linking immunoprecipitation (CLIP) studies demonstrating an enrichment of RBP-binding sites toward the very ends of 3'-UTRs (95, 136). Interestingly, increased accessibility correlates with increased sequence conservation in the last few hundred nucleotides toward the ends of alternative 3'-UTR isoforms (**Figure 1b**) (I. Singh & C. Mayr, unpublished data). The general association between RNA sequence conservation and accessibility in 3'-UTRs is reminiscent of what was also found in introns and ribozymes (109, 122). The generation of alternative 3'-UTRs seems to enable increased accessibility of 3'-UTR regulatory elements that are otherwise buried within long 3'-UTRs.

In addition to facilitating the accessibility of regulatory elements, alternative 3'-UTRs can also differentially regulate gene expression. Most genes that generate alternative 3'-UTR isoforms have evolved ways to regulate 3'-UTR ratios (81). Embryonic tissues tend to express shorter 3'-UTRs than differentiated tissues (63, 73, 115, 130). Across differentiated tissues, alternative 3'-UTRs are usually expressed in a cell type-specific manner (**Figure 1c**) (33, 81) and can change through activation of signaling pathways (32, 106) or disease (83, 87, 125). Usage of the most proximal polyadenylation (pA) site in these 3'-UTRs results in mRNA transcripts with rather short 3'-UTRs (median length of 400 nt). These are similar in length to mRNA transcripts generating only one 3'-UTR isoform (median length of 505 nt). This indicates that usage of proximal pA sites can result in the loss of the majority of regulatory elements present in longer 3'-UTRs.

### BIOLOGICAL FUNCTIONS OF 3'-UTRs

3'-UTRs are best known to determine the fate of mRNAs through the regulation of mRNA stability, translation, and mRNA localization. 3'-UTR functions are mediated by RBPs that bind to 3'-UTRs in a sequence- or structure-dependent manner. However, RBPs act only as adaptors that connect 3'-UTRs to effector proteins. Thus, the biological consequences that are mediated by 3'-UTRs depend on the functions of the effector proteins. For example, mRNA destabilization is achieved through RBPs that recruit a deadenylase (**Figure 2a**) (147), whereas translational

repression is accomplished by RBPs that recruit decapping enzymes (22). 3'-UTRs also determine the fate of newly translated proteins. This is facilitated by RBPs that recruit effector proteins that are transferred in a cotranslational manner onto the nascent peptide chain (9) (**Figure 2b**). This indicates that 3'-UTRs can mediate PPIs, and thus, regulate both protein localization and protein function (9).

### 3'-UTRs Mediate Protein-Protein Interactions


During translation of mRNAs, the nascent peptide chain and the 3'-UTR come into close proximity which seems to enable the transfer of proteins from the vicinity of the 3'-UTR onto the nascent peptide chain. Recruitment of protein interaction partners (PIPs) to the nascent peptide chain by 3'-UTRs may increase the association rate of the interacting proteins, and thus, may enable 3'UTR-dependent PPIs.

**3'-UTRs can regulate protein localization independently of mRNA localization.** It is commonly thought that the information for the formation of PPIs is fully contained within the interacting proteins themselves. This is illustrated in experiments studying PPIs by the use of constructs that contain only the coding regions (54). It is also generally assumed that proteins find their binding partners through diffusion and random collision. However, Berkovits & Mayr (9) made an observation that is not consistent with this prevalent model through their discovery that the 3'-UTR of *CD47* is required for the PPI between CD47 and SET. CD47 is known as a plasma membrane receptor that acts as a don't-eat-me signal (55), and the *CD47* gene uses APA to produce mRNA transcripts with either a short or a long 3'-UTR. Berkovits & Mayr called the protein that was generated from the short *CD47* 3'-UTR isoform CD47-SU and called the protein that was made from the long 3'-UTR isoform CD47-LU. Although CD47-SU and CD47-LU have identical amino acid sequences, Berkovits & Mayr demonstrated that, of the two, only CD47-LU interacts with SET (**Figure 2c**). This indicated that the formation of the PPI between CD47-LU and SET required recruitment of SET by the long *CD47* 3'-UTR (9). This PPI has functional consequences, as it promotes plasma membrane localization of CD47-LU, whereas CD47-SU is predominantly expressed intracellularly.

Importantly, Berkovits & Mayr demonstrated that the difference in CD47 protein localization was not due to differences in mRNA localization. Instead, a model was proposed and experimentally verified in which the RBP HuR binds only to the long, not the short, *CD47* 3'-UTR. HuR binding was responsible for the recruitment of SET to the long 3'-UTR (9, 13). During translation of the CD47 polypeptide chain, SET seems to be transferred onto the newly made CD47-LU protein, where SET binds to positively charged amino acids in the cytoplasmic domains of CD47-LU. SET also interacts with RAC1, and active RAC1 mediates efficient surface localization of CD47-LU (9, 124). In contrast, SET is not recruited to the short *CD47* 3'-UTR and does not interact with CD47-SU, leading to inefficient localization of CD47-SU to the plasma membrane. Thus, CD47-LU largely represents the well-known surface CD47 protein (55).

The study also expanded these findings to show that long 3'-UTRs of genes encoding other plasma membrane proteins, such as CD44, Integrin  $\alpha$ 1, and the BAFF receptor, were also able to increase cell surface expression of their cognate proteins. These candidates were investigated because they had the minimal requirements necessary for 3'-UTR-dependent protein localization. These requirements included HuR-binding sites in their long 3'-UTRs that facilitate SET recruitment and positively charged amino acids in their cytoplasmic protein domains necessary for SET binding (9). Therefore, 3'-UTRs can mediate PPIs, and thus they enable subcellular localization of proteins independently of mRNA localization.

**3'-UTR-mediated prerecruitment of the signal recognition particle to the ribosome.** 3'-UTR-mediated recruitment of proteins to the nascent peptide chain was recently confirmed in yeast using ribosome profiling of ribosome subpopulations. The 3'-UTRs of membrane protein-encoding genes, but not the 3'-UTR of a gene encoding a cytosolic protein, can recruit the signal recognition particle (SRP) to the ribosome before the polypeptide chain encoding the signal sequence has become exposed (19). The prevailing model of SRP recruitment to ribosomes has



been that translation of mRNAs encoding secretory proteins begins in the cytoplasm and, following exposure of the hydrophobic signal sequence at the exit channel of the ribosome, SRP binds to the signal sequence. This leads to halted translation and transfer of the SRP-bound ribosome to the surface of the endoplasmic reticulum. Once the ribosome is localized to the endoplasmic reticulum, translation resumes and the membrane protein is folded into the membrane (135). However, using co-immunoprecipitation (co-IP) of SRP72 followed by ribosome profiling, Chartron and colleagues (19) showed that this model is only true for a minority of secretory proteins. For the vast majority of proteins, SRP is prerecruited to ribosomes in a 3'-UTR-dependent manner before the signal sequence has been translated and before the nascent peptide chain has become exposed (19). These findings demonstrate that the selective interaction of SRP with ribosomes that translate secretory proteins is for the majority of cases not contained within the hydrophobic amino acids of the nascent peptides, but rather is mediated by elements in the 3'-UTRs of the mRNAs that are being translated. This suggests that specific 3'-UTRs recruit to the site of translation proteins that subsequently bind the nascent peptide chain.

**Widespread occurrence of cotranslational cognate mRNA–protein complexes.** For both SET and SRP, which are transferred onto the nascent peptide chain in a 3'-UTR-mediated manner, the model contains a complex that consists of a translating ribosome, an mRNA that encodes the nascent protein, and the nascent protein with an mRNA-recruited cotranslational PIP (**Figure 2b**). Such complexes seem to be widespread in yeast. For example, purification of the SET1 histone methyltransferase complex revealed an enrichment of *SET1* mRNA. *SET1* mRNA copurified with four of the eight proteins of the SET1 histone methyltransferase complex in a process that was dependent on cytoplasmic expression of the mRNA and on translation (45). Therefore, it was concluded that a SET1 subcomplex assembles on the nascent SET1 protein in a cotranslational manner. Using RNA-IP on proteins that do not contain RNA-binding domains, Duncan & Mata (31) showed that such cotranslational complexes occur in 38% of tested cases (**Figure 2b**). These data strongly suggest that the protein used for RNA-IP binds to the nascent polypeptide chain and that cotranslational PPIs that involve the corresponding mRNAs are widespread.

---

## Figure 2

Models of 3'-UTR-mediated functions. (a) RBPs (*red, orange*) bound to 3'-UTRs recruit effector proteins (*blue*) that determine the fate of mRNAs. (b) RBPs (*red, orange*) bound to 3'-UTRs recruit effector proteins (*blue*) that are transferred onto the nascent peptide chain (*dark green*), resulting in a 3'-UTR-dependent protein complex. (c) Co-IP of endogenous SET using GFP-trap after transfection of GFP-CD47-SU or GFP-CD47-LU in HEK293 cells. Two percent of input was loaded. (d) Model of alternative PIPs recruited by alternative 3'-UTRs despite the sequence of the short 3'-UTR being fully contained within the long 3'-UTR. It is possible that the mRNA structure occludes the regulatory elements in the short 3'-UTR within the context of the long 3'-UTR. Effector proteins are shown in light blue, dark blue, and purple. (e) Model of *peripherin* mRNA clustering by an RBP that binds to the *peripherin* 3'-UTR and clusters the mRNAs through multimerization. This brings nascent peripherin proteins together and may allow them to oligomerize using microtubules (*blue*) as scaffold. (f) In bacterial operons, genes encoding subunits of protein complexes are in close proximity and are translated simultaneously. This facilitates protein complex formation. In evolved operons in eukaryotes, the 3'-UTR recruits various PIPs to the site of translation. This has the advantage of being independent of simultaneous translation of the PIPs and allows the recruitment of different PIPs that enable formation of substoichiometric protein complexes with diverse compositions. (g) mRNA localization through a microtubule motor protein (*purple*) that connects the 3'-UTR to cytoskeletal filaments using an RBP (*red*) as adaptor. (h) An actin-binding domain in the N terminus of Abp140 binds to actin filaments and transports the cognate mRNA using actin retrograde flow. Abbreviations: 3'-UTRs, 3'-untranslated regions; co-IP, co-immunoprecipitation; GFP, green fluorescent protein; IgG, immunoglobulin G; LU, long 3'-untranslated region; PIP, protein interaction partner; RBP, RNA-binding protein; SU, short 3'-untranslated region; WB, Western blotting. Panel *c* adapted from Reference 9.

**Alternative 3'-UTRs mediate the multifunctionality of proteins.** In the case of CD47, the alternative 3'-UTRs not only regulate protein localization but also determine alternative functions of CD47. When cells that expressed either CD47-LU or CD47-SU were exposed to  $\gamma$ -irradiation, the cells that expressed CD47-LU survived whereas the cells that expressed CD47-SU died (9). The functional difference between CD47-SU and CD47-LU was due to alternative 3'-UTR-dependent PIPs (**Figure 2d**). As discussed earlier, SET binds to CD47-LU, but not to CD47-SU. Consequently, the binding domains in CD47-SU are available to interact with proteins other than SET. The CD47-SU-specific PIPs were shown to be responsible for cell death induction after  $\gamma$ -irradiation (85). These findings demonstrate that the information that determines protein complex formation as well as protein function is not always contained within the amino acid sequence; it can be encoded in the noncoding parts of mRNAs, specifically within 3'-UTRs.

**3'-UTR-mediated cotranslational complexes may facilitate homopolymerization of filaments.** Taking these new findings into account, earlier experimental evidence can be reinterpreted. For example, Chang and colleagues (16) studied the assembly of intermediate filaments in mammalian cells using peripherin as a model. Like the previously mentioned yeast studies, they identified cotranslational cognate mRNA–protein complexes together with a clustering of *peripherin* mRNAs. Interestingly, they showed that mRNA clustering depends on the presence of the *peripherin* 3'-UTR, translation of peripherin protein, and polymerized microtubules. Thus, a model can be proposed in which mRNA clustering facilitates cotranslational homopolymerization of filaments (**Figure 2e**). mRNA clustering can be achieved through an RBP that binds and connects different *peripherin* 3'-UTRs. This brings the ribosomes translating *peripherin* mRNAs into proximity with one another so that nascent peripherin subunits can assemble on microtubules. Such a model is consistent with early reports on the association of nascent peptides, including vimentin and myosin, with the cytoskeleton (36). As vimentin, myosin, and peripherin are built through homotypic coiled-coiled interactions, it is possible that polymerization of intermediate filament chains takes place cotranslationally and involves the coordinated synthesis of mRNAs that closely associate through their 3'-UTRs.

**3'-UTR-mediated protein–protein interactions: open questions and future directions.** 3'-UTR-mediated cotranslational protein transfer is a conserved and widespread mechanism in both yeast and human cells. An emerging picture from the studies described above is that 3'-UTRs can assist cotranslational multiprotein complex assembly, protein oligomerization, and the formation of PPIs (9, 16, 19, 31, 45). These data suggest that although PPIs take place between two proteins, the initiation of binding is facilitated through recruitment of a PIP by a 3'-UTR. However, the number of edges of the cotranslational mRNA–protein network seems to be larger in human cells than in yeast. In yeast, mRNAs recruit only a few mRNA-dependent PIPs, and the recruited PIPs generally interact with only one to three mRNAs (31, 45). By contrast, in human cells dozens or even hundreds of 3'-UTR-dependent PIPs have been found (79, 85).

The recruitment of many different 3'-UTR-dependent PIPs may enable the multifunctionality of proteins, even in the absence of alternative 3'-UTRs. For example, a single 3'-UTR could assist in the cotranslational formation of several substoichiometric protein complexes or subcomplexes. For processes that require the presence of several proteins in a specific location, the assumption that proteins are able to freely diffuse makes it unlikely that all of them are present simultaneously at the required location. However, through 3'-UTR-dependent formation of subcomplexes that contain the majority of these factors, only one protein localization event would be necessary. This suggests that 3'-UTRs may preorganize ligand binding sites of a protein complex. Preorganization reduces the freedom for binding events of individual components, which, in turn, decreases the number



of nonproductive configurations and reduces entropy. Such 3'-UTR-dependent preorganization may even increase reaction rates, as it reduces the freedom of motion of individual proteins (141). Thus, 3'-UTR-dependent PPIs enable cooperativity.

The topology of the cotranslational complexes that enable 3'-UTR-dependent PPIs is very reminiscent of translational operons in bacteria. Operons organize genes that encode distinct subunits of protein complexes to facilitate cotranslational protein complex assembly (116, 140). Cotranslational 3'-UTR-dependent protein complex assembly can be viewed as an evolved eukaryotic operon because the recruitment of PIPs by 3'-UTRs has several key advantages: A lack of dependency on simultaneous translation of the PIPs makes it more flexible, and the ability to recruit several different PIPs enables the formation of 3'-UTR-dependent protein complexes with diverse compositions (**Figure 2f**).

### 3'-UTRs Regulate mRNA Localization

Although it was initially thought that mRNA localization happens only in the case of a few select mRNAs in polarized cells, it seems that mRNA localization is actually very widespread. Large-scale localization experiments in *Drosophila* embryos showed that 71% of mRNAs have a nonuniform distribution within cells (77).

**mRNA localization through association with motor proteins.** Localization of mRNAs is usually mediated by localization signals in 3'-UTRs that are bound by RBPs. These RBPs in turn bind motor proteins, which result in movement of the mRNAs along filaments of the cytoskeleton (**Figure 2g**). One of the best-characterized examples of this is the asymmetric distribution of *ASH1* mRNA to the bud tip during cell division in yeast, which is required for the suppression of mating type switching in the daughter cells (98). *ASH1* mRNA contains four *cis*-elements, including one located in the 3'-UTR, which are required for proper localization. These elements are bound by two RBPs, She2p and She3p. She3p also binds to a myosin motor protein, Myo4p, which transports the mRNA–protein complex along actin microfilaments to the bud tip.

Other mRNAs, including *oskar* mRNA in *Drosophila* oocytes, are transported along microtubules. Local function of Oskar is required to promote the localization of *nanos* mRNA, which is translated in the embryo and forms a gradient important for proper embryo patterning (82). *oskar* mRNA enters the oocyte from the nurse cells, a step that requires a 3'-UTR element and the microtubule motor protein dynein (57). Dynein is connected to the 3'-UTR through the RBPs Egl and BicD, which serve as adaptors (29). After oocyte entry, *oskar* mRNA uses a different 3'-UTR element, bound by Tropomyosin1-I/C, which connects *oskar* mRNA to kinesin-1, a microtubule motor protein that moves toward the plus end. This results in localization of *oskar* mRNA to the posterior pole of the oocyte (37, 56, 152). 3'-UTR-dependent recruitment of motor proteins, which is mediated by RBPs, has been described for the majority of localized mRNAs, including *CAMK2A*, *bicoid*, and  $\beta$ -*actin* (82, 91, 112).

**Additional mechanisms of mRNA localization.** In some cases, RBPs that bind to mRNAs do not directly interact with motor proteins, but they are connected through additional RBPs that serve as adaptors, such as BicD or She3p (98). Interestingly, such an adaptor can also be an early endosome, as was shown for RRM4 in a fungus (8, 70). Another unconventional mRNA transport mechanism was described for Abp140, which localizes to the distal pole in the yeast mother cell. It forms a ternary complex consisting of the ribosome, the mRNA, and the nascent polypeptide chain. The actin-binding domain located in the N terminus of the nascent peptide chain binds to actin microfilaments and uses actin retrograde transport for localization (**Figure 2b**) (67).

**mRNA association with membranes and the cytoskeleton.** mRNAs and their bound RBPs often seem to be associated with cytoskeletal filaments or membranes (30, 59). They associate with cytoskeletal filaments during nucleocytoplasmic transport, but also while mediating their functions in the cytoplasm. For instance, the RBPs HuD and RAVR1 were shown to interact with actin-binding proteins and with microtubule-associated proteins, respectively (35, 52). mRNAs also bind to several components of vesicle coats, including COPI and COPII vesicle coat complexes (127, 146), and were found to localize to the surface of different organelles, such as mitochondria, endosomes, or the endoplasmic reticulum (25, 38, 60, 105). These examples show an intimate association between mRNAs, membranes, and the cytoskeleton. Intriguingly, repression by siRNAs or miRNAs also seems to happen on the surface of the endoplasmic reticulum (120). However, our understanding of the interactions among mRNAs, membranes, and the cytoskeleton is still rudimentary. The association of subgroups of mRNAs through their RBPs with different parts of the cytoskeleton may not only help to organize their localization but also enable compartmentalization of 3'-UTR functions (40).

### 3'-UTRs Play a Role as Independent Information Units

Most often, mRNAs are viewed as the templates for protein synthesis. However, mRNAs or parts of mRNAs, including cleaved fragments of 3'-UTRs, can also act as regulatory RNAs.

**mRNAs act like long noncoding RNAs.** For some of the well-studied model mRNAs, it was shown that the mRNA itself carries out a specific function that is independent of the encoded protein. This indicates that mRNAs can act as independent information units with functions similar to long noncoding RNAs. This was first demonstrated for *VegT* mRNA, which localizes to the vegetal pole in *Xenopus* oocytes (47). Its deletion impaired anchoring of other vegetal pole-associated mRNAs, including *BicC* and *Wnt11* mRNAs. This function was specific to *VegT* mRNA, as inhibition of *VegT* translation did not have such an effect. A protein-independent role was also found for *oskar* mRNA (61). *oskar* mRNA, but not Oskar protein, is required for the completion of oogenesis, as deletion of *oskar* mRNA results in an eggless phenotype (61). Importantly, expression of the *oskar* 3'-UTR was sufficient to rescue this phenotype, but the mechanism of action is currently unknown. It was speculated that *oskar* mRNA may localize or sequester an RBP. Alternatively, it may play a structural role, as was suggested for *VegT* mRNA (47, 61).

**Expression of 3'-UTRs independent from the coding region.** While analyzing CAGE libraries from human, mouse, and fly tissues, Mercer et al. (90) found 3'-UTRs that contained 5' caps. Although this was surprising, it was not a rare event: Thousands of genes have CAGE tags in their 3'-UTRs, including a subset of genes conserved between mouse and human transcripts (90). Several capped 3'-UTRs displayed discordant expression with the coding regions, which means that their expression is likely autonomous. Discordant expression was also seen in an independent study that analyzed polysome-derived mRNAs from mouse neurons (69). For several candidates, RNA-seq data revealed higher expression of the coding region than of the 3'-UTR, a finding likely explained by 3'-UTR shortening due to APA in this cell type. However, other candidates showed the opposite pattern, as they had substantially higher expression of the 3'-UTR compared to the coding region. *Sox11* was one of the most striking examples, with 6,000 reads mapping to the 3'-UTR and only eight reads mapping to the coding region. The expression difference of additional transcripts was less dramatic but still more than 20-fold (69). Differential expression of

a substantial number of candidates was confirmed using *in situ* hybridization and gave credence to the findings.

Dual function mRNAs were also recently described in bacteria, where stable 3' by-products of mRNA turnover act as small RNAs (sRNAs) (17). One of the sRNAs, CpxQ, was found to be produced from endonucleolytic cleavage by RNase E, which leaves a 5' monophosphate. The cleavage occurred a few nucleotides downstream of the stop codon and produced a 58-nt-long sRNA derived from the 3'-UTR (17). The 3'-UTR cleavage product is nearly completely conserved in sequence and acts as an Hfq-dependent repressor during the inner membrane stress response (17). Thus, a transcribed 3'-UTR can be converted into a sRNA through endonucleolytic cleavage.

Also in mammalian cell lines, researchers identified cleavage products with a 5' monophosphate that were generated by endonucleolytic cleavage. Transcriptome-wide degradome sequencing was initially performed to uncover the cleavage sites of the endonucleases of the miRNA pathway, *i.e.*, Ago2 and Drosha (64). Many cut sites were not dependent on these two endonucleases, but the enzymes responsible are currently unknown (10, 92, 108). Intriguingly, there is substantial overlap between the degradome and the 3'-UTR fragment data sets (64, 69, 90). Approximately 40% of the genes that were identified by degradome sequencing to harbor cut sites were found to generate capped 3'-UTR fragments (64, 90). This suggests that a subgroup of degradation products is *de novo* capped in the cytoplasm, an occurrence previously shown in mouse and human cell lines (94, 102). Current data suggest that the 5' fragments that are produced through endonucleolytic cleavage are rapidly degraded by the exosome and are usually not detectable (69, 108). However, the 3' fragments are protected at the 3' end by the poly(A) tail (90) and may be stabilized at the 5' end through a strong structural element like a pseudoknot (18), through the binding of an RBP, or when XRN1 levels are low (108).

### 3'-UTRs Regulate Protein Abundance

The first motifs discovered in 3'-UTRs were AU-rich elements leading to rapid mRNA decay (6, 21). AU-rich elements are preferentially found in genes whose expression requires tight regulation such as cytokines, proto-oncogenes, and immune-regulatory factors (6, 21). These genes are rather special, as their mRNA half-lives are shorter than 30 min, compared to the median half-life of the transcriptome which is 7 h (113). The importance of these regulatory elements in restricting protein expression is highlighted by the fact that their deletion is associated with cancer, chronic inflammation, and auto-immune disease (6).

Because of the drastic effects of AU-rich elements on the regulation of acute phase genes, it was suspected that 3'-UTRs may regulate protein abundance genome-wide. However, genome-wide analyses of 3'-UTR-mediated regulation of protein abundance generally showed widespread, modest changes (4, 111, 119). One explanation for the small effect may be that during unperturbed steady-state growth of cell lines, 3'-UTR elements do not substantially regulate protein abundance. However, in certain biological contexts, miRNAs and RBPs are important for the control of protein levels. Proto-oncogenes belong to the class of genes with short half-lives, and their expression levels are controlled by AU-rich elements and miRNA-binding sites (21). Removal of these elements through 3'-UTR shortening in cancer cell lines resulted in upregulation of their protein levels. This turned proto-oncogenes into oncogenes, as the increased protein expression was sufficient for oncogenic transformation (87).

In specific biological contexts, miRNAs and RBPs also degrade mRNA transcripts that do not belong to the class of short-lived acute phase proteins. In early animal development, gene expression is accomplished by maternal mRNAs. However, at the onset of zygotic transcription,

the maternal mRNAs need to be degraded. During zebrafish development, miR-430 is highly expressed at the onset of zygotic transcription and is the predominant regulator of maternal mRNA decay (41).

Another beautiful example of regulating protein abundance by RBPs occurs in the control of mitochondrial biogenesis after glucose depletion in yeast (128). When yeast cells are grown in high-glucose conditions, mitochondrial biogenesis is repressed; there is no need for oxidative phosphorylation because yeast cells prefer to obtain their energy from glycolysis (78). In *Saccharomyces cerevisiae*, nuclear-encoded mitochondrial transcripts are preferentially bound by the RBP Puf3 (38, 128). Under high-glucose conditions, Puf3 destabilizes the transcripts that encode mitochondrial proteins. However, glucose deprivation or growth on nonfermentable carbon sources induces mitochondrial biogenesis, which results in a switch from degradation to preferential translation of Puf3 target genes. The switch in Puf3 activity is mediated by phosphorylation induced by a lack of glucose (78). This example illustrates the RBP-mediated control of protein abundance upon environmental changes. It also illustrates how a single RBP can determine opposite outcomes for bound mRNAs. Such dual functions of RBPs have also been described for DDX6, HuR, and many others (137, 150).

### **THE 3'-UTR-SPECIFIC RNA-BINDING PROTEOME AT A GIVEN MOMENT DETERMINES THE FUNCTIONS OF A 3'-UTR**

The function of a given 3'-UTR regulatory element is influenced by both the RBP that directly binds to it and the RBPs that bind in its vicinity. This includes neighboring elements as well as distal elements that come into proximity through mRNA folding events. Thus, the functions of a 3'-UTR are determined by the ensemble of bound RBPs at a given moment. Notably, some RBPs can be loaded onto the mRNA in the nucleus, whereas others are added locally. Some RBPs are bound to 3'-UTRs with higher affinity, whereas others associate more transiently. To identify all RBPs bound to a single 3'-UTR, methods can be applied that were initially developed to determine the proteome associated with the long noncoding RNA Xist (23, 89). ChIRP-MS and RAP use biotinylated oligonucleotides to pull out endogenous RNAs followed by the identification of bound proteins by mass spectrometry. A similar method, called TRIP, was developed to examine the RBPs bound to mRNAs (84).

Fifteen years ago, the posttranscriptional operon theory was proposed by Keene & Tenenbaum (66). It stated that RBPs regulate subpopulations of mRNAs in a coherent manner and predicted that the mRNAs bound by an RBP would be functionally related. The best examples for RNA regulons are found in yeast, where, for example, Puf3 binds to a group of mRNAs that encode mitochondrial proteins (38, 128). To identify mRNAs that are bound by a single RBP, RNA-IP, and later CLIP (including HITS-CLIP, PAR-CLIP, iCLIP and eCLIP) were invented and widely applied (44, 71, 129, 132). CLIP uses an affinity tag or antibody against an RBP and precipitates the *in vivo* bound mRNAs after cross-linking. CLIP uncovered that the core motifs bound by RBPs are usually 3–7 nt long, are often degenerate, and are found in the majority of mRNAs (39). It also revealed that although many RBPs have hundreds or thousands of target mRNAs (76), a specific RBP usually binds only to 6–14% of its potential motifs *in vivo* (122, 132, 145).

### **Increased Specificity of RNA-Binding Proteins Through Cooperativity**

Considering that each RBP binds to thousands of mRNAs, it seems a daunting task to identify specific RBP–mRNA relationships that are functional in a given context. The fact that RBPs use

interactions with other proteins to increase the binding specificity to 3'-UTRs makes this process less overwhelming.

**Posttranslational modifications or protein interaction partners of RNA-binding proteins can change the binding specificity of RNA-binding proteins.** Campbell et al. (14) used an in vitro selection method called SEQRS, to identify the binding motifs of RBPs. They determined the motif of an RBP alone or when bound to a domain of a PIP. In both cases the core motif of the RBP was detected, but the sequence motif in the neighboring bases differed. This study demonstrated that the binding specificity of an RBP can be altered through association with a PIP, even if the PIP does not bind to the RNA. Similarly, the binding motif of an RBP can be altered through PTMs of the RBP. This was shown for HuR, which prefers binding to AU-rich sequences but shifts toward U-rich sequences upon phosphorylation (110).

**Ternary complex formation increases the specificity of mRNA-protein interactions.** RBPs often physically interact with other RBPs and were shown by CLIP studies to coassemble on mRNAs (93). The binding of two RBPs to neighboring sequence motifs in an mRNA has been shown in numerous examples to increase the binding specificity (1, 48). Such cooperativity was shown for the RBPs She2p and She3p when bound to *ASH1* mRNA. All binary interactions between She2p and *ASH1* mRNA, She3p and *ASH1* mRNA, and She2p and She3p resulted in low-affinity and low-specificity complexes (96). In contrast, ternary complex formation increased their affinity and enabled binding of the RBPs to *ASH1* mRNA with higher selectivity than to other hairpin mRNAs (96).

Another example of increased specificity through formation of a cooperative ternary complex was demonstrated by the crystal structure of the *Drosophila* RBPs Sxl and Unr and the *msl2* mRNA (49). Translational repression of *msl2* mRNA is required for sex chromosome dosage compensation in female flies. Sxl binds to U-rich sequences, and the RNA-binding motifs of Sxl and Unr are very abundant in the transcriptome. The extended RNA-binding motif in *msl2* mRNA contains 16 nt and is recognized by Sxl together with Unr. Cooperative protein complex formation resulted in a 1,000-fold increase in RNA binding affinity. Thus, this and other structures demonstrated how general and widespread mRNA motifs can be specifically recognized by cooperative ternary complex formation (49, 80).

### **Binding of RNA and Proteins in the Nucleus Can Influence Cytoplasmic 3'-UTR Functions**

Although most 3'-UTR functions are carried out in the cytoplasm, some RBPs bind to 3'-UTRs in the nucleus (118). For example, it is important to recruit the RBP She2p to *ASH1* mRNA in the nucleus, as mutant She2p that is unable to enter the nucleus results in impaired cytoplasmic localization of *ASH1* mRNA (98). Nuclear She2p is also necessary for the nuclear recruitment of the translational repressor Puf6p to *ASH1* mRNA, although it acts in the cytoplasm (114). Furthermore, nuclear events are important for *oskar* mRNA localization, as deposition of the exon junction complex during splicing is required for cytoplasmic localization of *oskar* mRNA (43).

Additional nuclear events are mediated through cotranscriptional loading of RBPs at promoters and seem to play a general role in the regulation of cytoplasmic processes of mRNAs. In yeast, promoters control mRNA stability, translation, and mRNA localization (12, 126, 151). Although glucose starvation leads to the translational repression of most yeast mRNAs, a subset remains actively translated. Binding of HSF1 to promoters determines the group of mRNAs that is being translated during glucose starvation. HSF1 binding may result in the recruitment of an RBP that

is exported with the mRNAs and specifies translation in the cytoplasm (151). A similar mechanism was proposed for the influence of promoters on mRNA stability, where swapping the upstream activating sequence (UAS) of a promoter changed the mRNA decay rate of the transcript. As changing the UAS did not alter the mRNA sequence, promoter-mediated recruitment of specific factors that influence cytoplasmic processes seems likely (12). Such promoter-mediated recruitment of an RBP was recently demonstrated in *Drosophila*. Loading of HuR at specific promoters influences the recognition of alternative pA sites (101). Thus, RBPs recruited to promoters can change whether regulatory elements are functional in 3'-UTRs.

### Local mRNA Functions Are Enabled Through Local Binding of RNA-Binding Proteins to 3'-UTRs

Local recruitment of RBPs to 3'-UTRs can also change 3'-UTR functions. One of the classical examples for mRNA transport in neurons is the localization of  $\beta$ -actin mRNA from the soma to the synapse. The zipcode binding protein IGF2BP1 (ZBP1, IMP-1) binds to a localization element in the 3'-UTR of  $\beta$ -actin mRNA, which is necessary for mRNA localization, but  $\beta$ -actin mRNA is released from IGF2BP1 binding after reaching the synapse. This event is triggered by phosphorylation of IGF2BP1, as only unphosphorylated Y396 can bind  $\beta$ -actin mRNA. Phosphorylation is mediated by Src kinase, whose expression is locally restricted to the synapse. During transport,  $\beta$ -actin mRNA is translationally repressed, but phosphorylation of IGF2BP1 also leads to translational activation of  $\beta$ -actin mRNA (53). Thus, the new local environment, characterized by expression of Src kinase, defines the end point of  $\beta$ -actin mRNA localization and enables its local functions. PTMs that occur in a spatially or temporally restricted manner are widely used to achieve dynamic changes in RBP composition at 3'-UTRs and, thus, in 3'-UTR function. For example, in certain stress conditions, HuR is ubiquitinated through addition of a K29 chain, a modification that does not induce degradation of HuR, but rather releases bound HuR from several of its target mRNAs. The release of HuR changes the fate of the mRNAs from stabilization to destabilization (150).

In addition to local expression of enzymes that add PTMs, local expression of RBPs can also change 3'-UTR functions. For example, *nanos* mRNA localizes to the posterior pole of the *Drosophila* embryo and establishes a morphogen gradient (82). However, it does not appear to be actively transported to the posterior pole because *nanos* mRNA is found throughout the embryo. Also throughout the embryo, the RBP Smaug binds to the *nanos* 3'-UTR and recruits a deadenylase, resulting in *nanos* mRNA decay. However, at the posterior pole, it was suggested that the RBP Oskar replaces Smaug, preventing Smaug from binding to the *nanos* 3'-UTR, which results in active translation and accumulation of Nanos protein at the posterior pole (147). This shows how the binding of locally restricted *trans*-acting factors to 3'-UTRs can determine local functions of mRNAs and their encoded proteins.

### Each Alternative 3'-UTR Isoform Has a Specific RNA-Binding Protein Composition and, Thus, Function

Differentiation and alterations in the cellular environment are associated with changes in signaling pathways that are able to change RBP composition at 3'-UTRs both in *cis* and in *trans*. Changes in the PTMs of RBPs can alter the association of an RBP with a 3'-UTR and, thus, alter the function of a 3'-UTR (53, 110, 150) (see section titled Posttranslational modifications or protein interaction partners of RNA-binding proteins can change the binding specificity of RNA-binding proteins). Furthermore, signaling can also change 3'-UTR ratios (32, 63, 81, 87, 106,

125). This can result in a switch in protein localization, protein abundance, and protein function, which has been shown for BDNF, for the oncogene IMP-1, and for CD47, respectively (3, 9, 75, 87).

**Functional discrepancy of 3'-UTR regulatory elements when analyzed in isolation or within a larger sequence context.** APA results in the expression of mRNAs with either short or long 3'-UTRs. The sequence of shorter 3'-UTRs is fully contained in longer 3'-UTRs (**Figure 1c**), therefore, it is commonly expected that an RBP that binds to the shorter 3'-UTR should also bind to the longer 3'-UTR. However, this is not necessarily the case, as the longer 3'-UTRs form their own distinct secondary and tertiary RNA structures that may occlude binding sites that are accessible in the context of shorter 3'-UTRs. Although this relationship has not been conclusively established for alternative 3'-UTRs, it was demonstrated that the functional effects of expressing smaller 3'-UTR pieces led to substantially different protein abundances when compared to full-length 3'-UTRs (72). Whereas most smaller 3'-UTR pieces (50-mers to 400-mers) had activating effects on protein abundance, the two full-length 3'-UTRs (*PIMI*, approximately 1,300 nt; *Hmga2*, approximately 2,900 nt) that were examined were both highly repressive (72, 88).

Such a discrepancy was also seen when even smaller 3'-UTR regulatory elements (8-mers) were tested in isolation or within the context of a larger endogenous sequence (142). When newly identified functional 3'-UTR elements were tested as 8-mers with respect to the regulation of protein abundance, all of them were either activating or repressive. However, when they were placed within a 500-nt endogenous sequence context, a significant fraction had opposite effects and the great majority had no regulatory effects on protein abundance (142). A likely explanation for this phenomenon is that a new local secondary structure is formed within the larger sequence context that prevents the accessibility or changes the folding of the isolated 8-mers. It is also possible that most RBPs do not act alone but rather compete or cooperate with RBPs that bind adjacently (14, 62). Thus, within the new sequence context, cooperativity with another RBP may alter the functional outcome of the RBP. It is also possible that binding of the RBP alone may regulate protein abundance, but interaction with a binding partner may result in the recruitment of a 3'-UTR-dependent PIP. Thus, the element would still be functional, but it would mediate a function that was not interrogated by the assay.

The observations that functional RBP-binding sites can be silent or have opposing effects when put into a new sequence context have important implications for alternative 3'-UTRs. They suggest that each alternative 3'-UTR isoform will have its own repertoire of functional RBP-binding sites, even if the sequence of the shorter 3'-UTRs is fully contained within longer 3'-UTRs. Thus, in a specific cellular condition, shorter 3'-UTRs may be associated with different RBPs than their corresponding longer 3'-UTRs (**Figure 2d**) (79, 85). However, each 3'-UTR isoform likely can adopt several different configurations that depend on the activity of signaling pathways, expression of RBPs, their cofactors, their PTMs, and their subcellular localization (53, 110, 150). Taken together, it seems that longer 3'-UTRs in particular have a lot of latent regulatory capacity that is regulated through accessibility (122). Yet, our understanding of how accessibility of 3'-UTR regulatory elements is controlled is still rudimentary. 3'-UTR structure analyses will help to elucidate this, but, ideally, they need to be performed in vivo on full-length 3'-UTRs in different contexts.

**Different biophysical properties of short and long 3'-UTRs.** For instance, short 3'-UTRs are predominantly found in the cytoplasm, whereas long 3'-UTRs can be part of liquid droplets and hydrogels (46, 65). The reasons for the difference in distribution are currently not known. It

is possible that long 3'-UTRs bind more RBPs than short 3'-UTRs. Because RBPs are enriched in protein domains with an overabundance of a few amino acids, called low-complexity (LC) domains (15), they are often intrinsically disordered. This causes most of their residues to be accessible and, thus, these residues represent major interacting domains for other proteins (40, 138). This feature enables RBPs to function as network hubs, as they were shown to have, on average, more PIPs than other protein classes (5). The multivalency of RBPs together with their LC domains allows liquid-liquid phase transitions of messenger ribonucleoproteins (65, 138). Phase transitions seem to play a very important role in the regulation of RNA-protein interactions, as they enable an increase in the local concentration and compartmentalization of factors without the need for membranes, providing a unique local environment that is distinct from the cytoplasm (46, 65, 134). Thus, in addition to a specialized local environment found on the surface of organelles (131), localization to liquid droplets can also change the range of RBPs that are available for binding to 3'-UTRs. There also seems to be interdependency between the mRNA and protein components within the droplet. On the one hand, as mRNAs are differentially bound by RBPs, mRNAs are able to determine the protein composition and the biophysical properties of the droplet (148). On the other hand, the biophysical properties of the droplet determine which RNAs are retained in the droplet (99). As liquid droplet formation may serve as a dynamic compartmentalization tool to enable local mRNA processes, 3'-UTR length itself may promote localized processes and may contribute to compartmentalization of cellular functions (133).

## NEW GLOBAL APPROACHES IDENTIFY NOVEL RNA-BINDING PROTEIN MOTIFS AND FUNCTIONAL 3'-UTR ELEMENTS

In yeast, Puf3 preferentially binds to mRNA transcripts that encode mitochondrial proteins (38, 128). However, binding of Puf3 can result in translational repression or activation (78). This indicates that it is not the sequence element per se, but rather the status (PTMs, PIPs) of the *trans*-acting factor, that determines the functional outcome. This interpretation is supported by recent studies that used fluorescence-based screens to identify new functional elements in 3'-UTRs (100, 142, 149). All of the screens used small 3'-UTR fragments to search for 3'-UTR elements that regulate protein abundance and found similar numbers of activating and repressive elements. However, interestingly, when the functional 3'-UTR fragments were placed within a larger endogenous sequence context, a large fraction showed either no effect or the opposite effect on protein abundance (142). This suggests that functional screens that use small 3'-UTR fragments can identify elements with a potential for regulation, but they are not able to identify the biological context or endogenous 3'-UTR that will produce a specific phenotype when harboring the element. Therefore, such screens are able to catalog RBP-binding sites with potential function and are comparable to bioinformatic predictions of miRNA targets (2). Further experiments are necessary to examine if a specific binding site is functional within a particular 3'-UTR as well as within a certain biological context.

The Giraldez laboratory (144) recently developed a screen that may go a step further in identifying the functional 3'-UTR elements that regulate mRNA stability in a specific biological context. The RNA-element selection assay (RESA) uses random fragmentation of 3'-UTR sequences to detect regions that contain stabilizing and destabilizing elements. For a proof-of-principle analysis, it was used to identify functional 3'-UTR elements at the onset of zygotic gene expression during zebrafish development. At this stage, all maternal mRNAs need to be degraded, and RESA confirmed that this is predominantly accomplished by miR-430 (144). This novel approach may also identify functional RBP regulons in different biological contexts. The combination of such



functional screens with newly developed in vitro methods that identify high-affinity motifs of RBPs (14, 74, 104) will help to shed light on the multilayered regulation accomplished by 3'-UTRs.

## CURRENT ISSUES AND HOW THEY CAN BE OVERCOME

It is currently not known how many RBPs bind to a specific 3'-UTR or to a pair of alternative 3'-UTRs. If such data were to exist for many 3'-UTRs, it would tremendously increase our understanding of 3'-UTR biology. It is also important to learn about the dynamics of RBP binding to specific 3'-UTRs in steady-state or altered cellular conditions. This could be accomplished by performing ChIRP-MS, RAP, or TRIP (23, 84, 89) on many 3'-UTRs in different cellular conditions.

Many RBPs bind to U-rich regions. Assuming that this observation is not a CLIP-generated artifact (121), there is a need to develop assays to examine the in vivo binding sites and functions of these RBPs while they are bound to their PIPs (14). This would enable the integration of cooperative binding events with functional assays.

RBPs mediate 3'-UTR functions, but they are only the adaptors that connect the 3'-UTR *cis*-regulatory elements to effector proteins. The effector proteins are the true mediators of a specific phenotype. As the effector proteins are generally PIPs of RBPs, there is a need to comprehensively identify the protein interactome of RBPs (11).

Researchers who set out to globally examine 3'-UTR elements often used small 3'-UTR fragments to probe for functional elements (100, 142, 144, 149). When the effects of 3'-UTR fragments on protein output were compared with the effects of full-length 3'-UTRs, substantially different outcomes were observed (72, 142) (see section titled Functional discrepancy of 3'-UTR regulatory elements when analyzed in isolation or within a larger sequence context). This indicates that 3'-UTR elements function in the context of the whole 3'-UTR or mRNA. Thus, methods need to be developed that identify functional regulatory elements within each alternative 3'-UTR isoform, as the presence of the motif often does not correlate with its functionality. Additionally, there is a need to develop functional assays that investigate global 3'-UTR functions that use full-length and alternative 3'-UTR isoforms as input.

Lastly, the majority of assays that assessed global 3'-UTR functions have used protein abundance as a functional readout. Although these readouts are both well established and highly reliable, regulation of protein abundance is only one of the many functions of 3'-UTRs. Thus, new readout systems are needed that are able to globally determine additional 3'-UTR functions such as mRNA localization (123) or 3'-UTR-mediated transfer of PIPs, which can be investigated by performing the experiment shown in **Figure 2c** (9, 79).

## DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

The author apologizes to all colleagues whose work could not be cited because of space constraints. She is grateful to all the members of the Mayr laboratory for helpful discussions and for critical reading of the manuscript. Funding for work on 3'-UTRs was obtained from the Innovator Award of the Damon Runyon-Rachleff Cancer Foundation and the Island Outreach Foundation, the Pershing Square Sohn Cancer Research Alliance, the Memorial Sloan Kettering core grant

P30-CA008748, the National Cancer Institute grant U01-CA164190, and the National Institutes of Health Director's Pioneer Award DPI-GM123454.

## LITERATURE CITED

1. Achsel T, Bagni C. 2016. Cooperativity in RNA-protein interactions: the complex is more than the sum of its partners. *Curr. Opin. Neurobiol.* 39:146–51
2. Agarwal V, Bell GW, Nam J-W, Bartel DP. 2015. Predicting effective microRNA target sites in mammalian mRNAs. *eLife* 4:e05005
3. An JJ, Gharami K, Liao GY, Woo NH, Lau AG, et al. 2008. Distinct role of long 3' UTR BDNF mRNA in spine morphology and synaptic plasticity in hippocampal neurons. *Cell* 134:175–87
4. Baek D, Villen J, Shin C, Camargo FD, Gygi SP, Bartel DP. 2008. The impact of microRNAs on protein output. *Nature* 455:64–71
5. Baltz AG, Munschauer M, Schwanhauser B, Vasile A, Murakawa Y, et al. 2012. The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts. *Mol. Cell* 46:674–90
6. Barreau C, Paillard L, Osborne HB. 2005. AU-rich elements and associated factors: Are there unifying principles? *Nucleic Acids Res.* 33:7138–50
7. Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* 136:215–33
8. Baumann S, Pohlmann T, Jungbluth M, Brachmann A, Feldbrügge M. 2012. Kinesin-3 and dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. *J. Cell Sci.* 125:2740–52
9. Berkovits BD, Mayr C. 2015. Alternative 3' UTRs act as scaffolds to regulate membrane protein localization. *Nature* 522:363–67
10. Boehm V, Haberman N, Ottens F, Ule J, Gehring NH. 2014. 3' UTR length and messenger ribonucleoprotein composition determine endocleavage efficiencies at termination codons. *Cell Rep.* 9:555–68
11. Brannan KW, Jin W, Huelga SC, Banks CA, Gilmore JM, et al. 2016. SONAR discovers RNA-binding proteins from analysis of large-scale protein-protein interactomes. *Mol. Cell* 64:282–93
12. Bregman A, Avraham-Kelbert M, Barkai O, Duek L, Guterman A, Choder M. 2011. Promoter elements regulate cytoplasmic mRNA decay. *Cell* 147:1473–83
13. Brennan CM, Gallouzi IE, Steitz JA. 2000. Protein ligands to HuR modulate its interaction with target mRNAs in vivo. *J. Cell Biol.* 151:1–14
14. Campbell ZT, Bhimsaria D, Valley CT, Rodriguez-Martinez JA, Menichelli E, et al. 2012. Cooperativity in RNA-protein interactions: global analysis of RNA binding specificity. *Cell Rep.* 1:570–81
15. Castello A, Fischer B, Eichelbaum K, Horos R, Beckmann BM, et al. 2012. Insights into RNA biology from an atlas of mammalian mRNA-binding proteins. *Cell* 149:1393–406
16. Chang L, Shav-Tal Y, Trcek T, Singer RH, Goldman RD. 2006. Assembling an intermediate filament network by dynamic cotranslation. *J. Cell Biol.* 172:747–58
17. Chao Y, Vogel J. 2016. A 3' UTR-derived small RNA provides the regulatory noncoding arm of the inner membrane stress response. *Mol. Cell* 61:352–63
18. Chapman EG, Costantino DA, Rabe JL, Moon SL, Wilusz J, et al. 2014. The structural basis of pathogenic subgenomic flavivirus RNA (sfRNA) production. *Science* 344:307–10
19. Chartron JW, Hunt KC, Frydman J. 2016. Cotranslational signal-independent SRP preloading during membrane targeting. *Nature* 536:224–28
20. Chen C-Y, Chen S-T, Juan H-F, Huang H-C. 2012. Lengthening of 3'UTR increases with morphological complexity in animal evolution. *Bioinformatics* 28:3178–81
21. Chen C-YA, Shyu A-B. 1995. AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem. Sci.* 20:465–70
22. Chen Y, Boland A, Kuzuoğlu-Öztürk D, Bawankar P, Loh B, et al. 2014. A DDX6-CNOT1 complex and W-binding pockets in CNOT9 reveal direct links between miRNA target recognition and silencing. *Mol. Cell* 54:737–50
23. Chu C, Zhang QC, da Rocha ST, Flynn RA, Bharadwaj M, et al. 2015. Systematic discovery of Xist RNA binding proteins. *Cell* 161:404–16
24. Crick FH. 1958. On protein synthesis. *Symp. Soc. Exp. Biol.* 12:138–63

25. Cui XA, Palazzo AF. 2014. Localization of mRNAs to the endoplasmic reticulum. *Wiley Interdiscip. Rev. RNA* 5:481–92
26. Dar D, Prasse D, Schmitz RA, Sorek R. 2016. Widespread formation of alternative 3' UTR isoforms via transcription termination in archaea. *Nat. Microbiol.* 1:16143
27. Dar D, Shamir M, Mellin JR, Koutero M, Stern-Ginossar N, et al. 2016. Term-seq reveals abundant ribo-regulation of antibiotics resistance in bacteria. *Science* 352:aad9822
28. Derti A, Garrett-Engle P, Macisaac KD, Stevens RC, Sriram S, et al. 2012. A quantitative atlas of polyadenylation in five mammals. *Genome Res.* 22:1173–83
29. Dienstbier M, Boehl F, Li X, Bullock SL. 2009. Egalitarian is a selective RNA-binding protein linking mRNA localization signals to the dynein motor. *Genes Dev.* 23:1546–58
30. Doller A, Schulz S, Pfeilschifter J, Eberhardt W. 2013. RNA-dependent association with myosin IIA promotes F-actin-guided trafficking of the *ELAV*-like protein HuR to polysomes. *Nucleic Acids Res.* 41:9152–67
31. Duncan CD, Mata J. 2011. Widespread cotranslational formation of protein complexes. *PLoS Genet.* 7:e1002398
32. Flavell SW, Kim T-K, Gray JM, Harmin DA, Hemberg M, et al. 2008. Genome-wide analysis of MEF2 transcriptional program reveals synaptic target genes and neuronal activity-dependent polyadenylation site selection. *Neuron* 60:1022–38
33. Floor SN, Doudna JA. 2016. Tunable protein synthesis by transcript isoforms in human cells. *eLife* 5:e10921
34. Friedman RC, Farh KK, Burge CB, Bartel DP. 2009. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 19:92–105
35. Fujiwara Y, Kasashima K, Saito K, Fukuda M, Fukao A, et al. 2011. Microtubule association of a neuronal RNA-binding protein HuD through its binding to the light chain of MAP1B. *Biochimie* 93:817–22
36. Fulton AB, L'Ecuyer T. 1993. Cotranslational assembly of some cytoskeletal proteins: implications and prospects. *J. Cell Sci.* 105:867–71
37. Gáspár I, Sysoev V, Komissarov A, Ephrussi A. 2017. An RNA-binding atypical tropomyosin recruits kinesin-1 dynamically to *oskar* mRNPs. *EMBO J.* 36:319–33
38. Gerber AP, Herschlag D, Brown PO. 2004. Extensive association of functionally and cytotopically related mRNAs with Puf family RNA-binding proteins in yeast. *PLoS Biol.* 2:E79
39. Gerstberger S, Hafner M, Tuschl T. 2014. A census of human RNA-binding proteins. *Nat. Rev. Genet.* 15:829–45
40. Gibson TJ. 2009. Cell regulation: determined to signal discrete cooperation. *Trends Biochem. Sci.* 34:471–82
41. Giraldez AJ, Mishima Y, Rihel J, Grocock RJ, Van Dongen S, et al. 2006. Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* 312:75–79
42. Grimson A, Farh KK-H, Johnston WK, Garrett-Engle P, Lim LP, Bartel DP. 2007. MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol. Cell* 27:91–105
43. Hachet O, Ephrussi A. 2004. Splicing of *oskar* RNA in the nucleus is coupled to its cytoplasmic localization. *Nature* 428:959–63
44. Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, et al. 2010. PAR-CLIP—a method to identify transcriptome-wide the binding sites of RNA binding proteins. *J. Vis. Exp.* 41:e2034
45. Halbach A, Zhang H, Wengi A, Jablonska Z, Gruber IM, et al. 2009. Cotranslational assembly of the yeast SET1C histone methyltransferase complex. *EMBO J.* 28:2959–70
46. Han TW, Kato M, Xie S, Wu LC, Mirzaei H, et al. 2012. Cell-free formation of RNA granules: Bound RNAs identify features and components of cellular assemblies. *Cell* 149:768–79
47. Heasman J, Wessely O, Langland R, Craig EJ, Kessler DS. 2001. Vegetal localization of maternal mRNAs is disrupted by VegT depletion. *Dev. Biol.* 240:377–86
48. Hennig J, Gebauer F, Sattler M. 2014. Breaking the protein–RNA recognition code. *Cell Cycle* 13:3619–20
49. Hennig J, Militti C, Popowicz GM, Wang I, Sonntag M, et al. 2014. Structural basis for the assembly of the Sxl–Unr translation regulatory complex. *Nature* 515:287–90

50. Hoffman Y, Bublik DR, Ugalde AP, Elkon R, Biniashvili T, et al. 2016. 3'UTR shortening potentiates microRNA-based repression of pro-differentiation genes in proliferating human cells. *PLoS Genet.* 12:e1005879
51. Hogan DJ, Riordan DP, Gerber AP, Herschlag D, Brown PO. 2008. Diverse RNA-binding proteins interact with functionally related sets of RNAs, suggesting an extensive regulatory system. *PLoS Biol.* 6:e255
52. Hüttelmaier S, Illenberger S, Grosheva I, Rüdiger M, Singer RH, Jockusch BM. 2001. Raver1, a dual compartment protein, is a ligand for PTB/hnRNPI and microfilament attachment proteins. *J. Cell Biol.* 155:775–86
53. Hüttelmaier S, Zenklusen D, Lederer M, Dichtenberg J, Lorenz M, et al. 2005. Spatial regulation of  $\beta$ -actin translation by Src-dependent phosphorylation of ZBP1. *Nature* 438:512–15
54. Huttlin EL, Ting L, Bruckner RJ, Gebreab F, Gygi MP, et al. 2015. The BioPlex network: a systematic exploration of the human interactome. *Cell* 162:425–40
55. Jaiswal S, Jamieson CH, Pang WW, Park CY, Chao MP, et al. 2009. CD47 is upregulated on circulating hematopoietic stem cells and leukemia cells to avoid phagocytosis. *Cell* 138:271–85
56. Jambor H, Brunel C, Ephrussi A. 2011. Dimerization of *oskar* 3' UTRs promotes hitchhiking for RNA localization in the *Drosophila* oocyte. *RNA* 17:2049–57
57. Jambor H, Mueller S, Bullock SL, Ephrussi A. 2014. A stem-loop structure directs *oskar* mRNA to microtubule minus ends. *RNA* 20:429–39
58. Jan CH, Friedman RC, Ruby JG, Bartel DP. 2011. Formation, regulation and evolution of *Caenorhabditis elegans* 3'UTRs. *Nature* 469:97–101
59. Jansen RP. 1999. RNA-cytoskeletal associations. *FASEB J.* 13:455–66
60. Jansen RP, Niessing D, Baumann S, Feldbrügge M. 2014. mRNA transport meets membrane traffic. *Trends Genet.* 30:408–17
61. Jenny A, Hachet O, Závorszky P, Cyrklaff A, Weston MDJ, et al. 2006. A translation-independent role of *oskar* RNA in early *Drosophila* oogenesis. *Development* 133:2827–33
62. Jens M, Rajewsky N. 2015. Competition between target sites of regulators shapes post-transcriptional gene regulation. *Nat. Rev. Genet.* 16:113–26
63. Ji Z, Tian B. 2009. Reprogramming of 3' untranslated regions of mRNAs by alternative polyadenylation in generation of pluripotent stem cells from different cell types. *PLoS ONE* 4:e8419
64. Karginov FV, Cheloufi S, Chong MM, Stark A, Smith AD, Hannon GJ. 2010. Diverse endonucleolytic cleavage sites in the mammalian transcriptome depend upon microRNAs, Drosha, and additional nucleases. *Mol. Cell* 38:781–88
65. Kato M, Han TW, Xie S, Shi K, Du X, et al. 2012. Cell-free formation of RNA granules: Low complexity sequence domains form dynamic fibers within hydrogels. *Cell* 149:753–67
66. Keene JD, Tenenbaum SA. 2002. Eukaryotic mRNPs may represent posttranscriptional operons. *Mol. Cell* 9:1161–67
67. Kilchert C, Spang A. 2011. Cotranslational transport of ABP140 mRNA to the distal pole of *S. cerevisiae*. *EMBO J.* 30:3567–80
68. Kim D, Kim J, Baek D. 2014. Global and local competition between exogenously introduced microRNAs and endogenously expressed microRNAs. *Mol. Cells* 37:412–17
69. Kocabas A, Duarte T, Kumar S, Hynes MA. 2015. Widespread differential expression of coding region and 3' UTR sequences in neurons and other tissues. *Neuron* 88:1149–56
70. König J, Baumann S, Koepke J, Pohlmann T, Zarnack K, Feldbrügge M. 2009. The fungal RNA-binding protein Rrm4 mediates long-distance transport of *ubi1* and *rbo3* mRNAs. *EMBO J.* 28:1855–66
71. König J, Zarnack K, Luscombe NM, Ule J. 2012. Protein–RNA interactions: new genomic technologies and perspectives. *Nat. Rev. Genet.* 13:77–83
72. Kristjánsson K, Fogarty EA, Grimson A. 2015. Systematic analysis of the *Hmg2* 3' UTR identifies many independent regulatory sequences and a novel interaction between distal sites. *RNA* 21:1346–60
73. Lackford B, Yao C, Charles GM, Weng L, Zheng X, et al. 2014. Fip1 regulates mRNA alternative polyadenylation to promote stem cell self-renewal. *EMBO J.* 33:878–89


74. Lambert N, Robertson A, Jangi M, McGeary S, Sharp PA, Burge CB. 2014. RNA Bind-n-Seq: quantitative assessment of the sequence and structural binding specificity of RNA binding proteins. *Mol. Cell* 54:887–900
75. Lau AG, Irier HA, Gu J, Tian D, Ku L, et al. 2010. Distinct 3'UTRs differentially regulate activity-dependent translation of brain-derived neurotrophic factor (BDNF). *PNAS* 107:15945–50
76. Lebedeva S, Jens M, Theil K, Schwanhäusser B, Selbach M, et al. 2011. Transcriptome-wide analysis of regulatory interactions of the RNA-binding protein HuR. *Mol. Cell* 43:340–52
77. Lecuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, et al. 2007. Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* 131:174–87
78. Lee CD, Tu BP. 2015. Glucose-regulated phosphorylation of the PUF protein Puf3 regulates the translational fate of its bound mRNAs and association with RNA granules. *Cell Rep.* 11:1638–50
79. Lee SH, Mayr C. 2017. *Alternative 3'UTRs regulate protein complex formation and post-translational modifications*. Presented at RNA Localization and Local Translation EMBO|EMBL Symp., Jul. 24, Barga, Italy
80. Leeper TC, Qu X, Lu C, Moore C, Varani G. 2010. Novel protein–protein contacts facilitate mRNA 3'-processing signal recognition by Rna15 and Hrp1. *J. Mol. Biol.* 401:334–49
81. Lianoglou S, Garg V, Yang JL, Leslie CS, Mayr C. 2013. Ubiquitously transcribed genes use alternative polyadenylation to achieve tissue-specific expression. *Genes Dev.* 27:2380–96
82. Martin KC, Ephrussi A. 2009. mRNA localization: gene expression in the spatial dimension. *Cell* 136:719–30
83. Masamha CP, Xia Z, Yang J, Albrecht TR, Li M, et al. 2014. CFIm25 links alternative polyadenylation to glioblastoma tumour suppression. *Nature* 510:412–16
84. Matia-González AM, Laing EE, Gerber AP. 2015. Conserved mRNA-binding proteomes in eukaryotic organisms. *Nat. Struct. Mol. Biol.* 22:1027–33
85. Mayr C. 2016. *3'UTR-mediated protein–protein interactions regulate protein functions*. Presented at The Complex Life of mRNA EMBO|EMBL Symp., Oct. 7, Heidelberg, Ger.
86. Mayr C. 2016. Evolution and biological roles of alternative 3'UTRs. *Trends Cell Biol.* 26:227–37
87. Mayr C, Bartel DP. 2009. Widespread shortening of 3'UTRs by alternative cleavage and polyadenylation activates oncogenes in cancer cells. *Cell* 138:673–84
88. Mayr C, Hemann MT, Bartel DP. 2007. Disrupting the pairing between *let-7* and *Hmga2* enhances oncogenic transformation. *Science* 315:1576–79
89. McHugh CA, Chen CK, Chow A, Surka CF, Tran C, et al. 2015. The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature* 521:232–36
90. Mercer TR, Wilhelm D, Dinger ME, Solda G, Korbie DJ, et al. 2011. Expression of distinct RNAs from 3' untranslated regions. *Nucleic Acids Res.* 39:2393–403
91. Miller S, Yasuda M, Coats JK, Jones Y, Martone ME, Mayford M. 2002. Disruption of dendritic translation of CaMKII $\alpha$  impairs stabilization of synaptic plasticity and memory consolidation. *Neuron* 36:507–19
92. Mino T, Murakawa Y, Fukao A, Vandenbon A, Wessels HH, et al. 2015. Regnase-1 and Roquin regulate a common element in inflammatory mRNAs by spatiotemporally distinct mechanisms. *Cell* 161:1058–73
93. Mitchell SF, Jain S, She M, Parker R. 2013. Global analysis of yeast mRNPs. *Nat. Struct. Mol. Biol.* 20:127–33
94. Mukherjee C, Bakthavachalu B, Schoenberg DR. 2014. The cytoplasmic capping complex assembles on adapter protein Nck1 bound to the proline-rich C-terminus of mammalian capping enzyme. *PLOS Biol.* 12:e1001933
95. Mukherjee N, Jacobs NC, Hafner M, Kennington EA, Nusbaum JD, et al. 2014. Global target mRNA specification and regulation by the RNA-binding protein ZFP36. *Genome Biol.* 15:R12
96. Muller M, Heym RG, Mayer A, Kramer K, Schmid M, et al. 2011. A cytoplasmic complex mediates specific mRNA recognition and localization in yeast. *PLOS Biol.* 9:e1000611
97. NCBI (Nat. Cent. Biotechnol. Inf.). 2016. *Genome database*. Bethesda, MD, updated Jul. 20. <https://www.ncbi.nlm.nih.gov/genome>
98. Niedner A, Edelman FT, Niessing D. 2014. Of social molecules: the interactive assembly of *ASH1* mRNA-transport complexes in yeast. *RNA Biol.* 11:998–1009

99. Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, et al. 2015. Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles. *Mol. Cell* 57:936–47
100. Oikonomou P, Goodarzi H, Tavazoie S. 2014. Systematic identification of regulatory elements in conserved 3' UTRs of human transcripts. *Cell Rep.* 7:281–92
101. Oktaba K, Zhang W, Lotz TS, Jun DJ, Lemke SB, et al. 2015. ELAV links paused Pol II to alternative polyadenylation in the *Drosophila* nervous system. *Mol. Cell* 57:341–48
102. Otsuka Y, Kedersha NL, Schoenberg DR. 2009. Identification of a cytoplasmic complex that adds a cap onto 5'-monophosphate RNA. *Mol. Cell. Biol.* 29:2155–67
103. Proudfoot NJ, Brownlee GG. 1976. 3' non-coding region sequences in eukaryotic messenger RNA. *Nature* 263:211–14
104. Ray D, Kazan H, Cook KB, Weirauch MT, Najafabadi HS, et al. 2013. A compendium of RNA-binding motifs for decoding gene regulation. *Nature* 499:172–77
105. Reid DW, Nicchitta CV. 2015. Diversity and selectivity in mRNA translation on the endoplasmic reticulum. *Nat. Rev. Mol. Cell Biol.* 16:221–31
106. Sandberg R, Neilson JR, Sarma A, Sharp PA, Burge CB. 2008. Proliferating cells express mRNAs with shortened 3' untranslated regions and fewer microRNA target sites. *Science* 320:1643–47
107. Schlackow M, Marguerat S, Proudfoot NJ, Bähler J, Erban R, Gullerova M. 2013. Genome-wide analysis of poly(A) site selection in *Schizosaccharomyces pombe*. *RNA* 19:1617–31
108. Schoenberg DR. 2011. Mechanisms of endonuclease-mediated mRNA decay. *Wiley Interdiscip. Rev. RNA* 2:582–600
109. Schultes EA, Spasic A, Mohanty U, Bartel DP. 2005. Compact and ordered collapse of randomly generated RNA sequences. *Nat. Struct. Mol. Biol.* 12:1130–36
110. Schulz S, Doller A, Pardini NR, Wilce JA, Pfeilschifter J, Eberhardt W. 2013. Domain-specific phosphomimetic mutation allows dissection of different protein kinase C (PKC) isotype-triggered activities of the RNA binding protein HuR. *Cell. Signal.* 25:2485–95
111. Selbach M, Schwanhausser B, Thierfelder N, Fang Z, Khanin R, Rajewsky N. 2008. Widespread changes in protein synthesis induced by microRNAs. *Nature* 455:58–63
112. Severt WL, Biber TU, Wu X, Hecht NB, DeLorenzo RJ, Jakoi ER. 1999. The suppression of testis-brain RNA binding protein and kinesin heavy chain disrupts mRNA sorting in dendrites. *J. Cell Sci.* 112:3691–702
113. Sharova LV, Sharov AA, Nedorezov T, Piao Y, Shaik N, Ko MS. 2009. Database for mRNA half-life of 19,977 genes obtained by DNA microarray analysis of pluripotent and differentiating mouse embryonic stem cells. *DNA Res.* 16:45–58
114. Shen Z, Paquin N, Forget A, Chartrand P. 2009. Nuclear shuttling of She2p couples *ASH1* mRNA localization to its translational repression by recruiting Loc1p and Puf6p. *Mol. Biol. Cell* 20:2265–75
115. Shepard PJ, Choi EA, Lu J, Flanagan LA, Hertel KJ, Shi Y. 2011. Complex and dynamic landscape of RNA polyadenylation revealed by PAS-Seq. *RNA* 17:761–72
116. Shieh YW, Minguez P, Bork P, Auburger JJ, Guilbride DL, et al. 2015. Operon structure and cotranslational subunit association direct protein assembly in bacteria. *Science* 350:678–80
117. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, et al. 2005. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res.* 15:1034–50
118. Singh G, Pratt G, Yeo GW, Moore MJ. 2015. The clothes make the mRNA: past and present trends in mRNP fashion. *Annu. Rev. Biochem.* 84:325–54
119. Spies N, Burge CB, Bartel DP. 2013. 3' UTR-isoform choice has limited influence on the stability and translational efficiency of most mRNAs in mouse fibroblasts. *Genome Res.* 23:2078–90
120. Stalder L, Heusermann W, Sokol L, Trojer D, Wirz J, et al. 2013. The rough endoplasmic reticulum is a central nucleation site of siRNA-mediated RNA silencing. *EMBO J.* 32:1115–27
121. Sugimoto Y, König J, Hussain S, Zupan B, Curk T, et al. 2012. Analysis of CLIP and iCLIP methods for nucleotide-resolution studies of protein–RNA interactions. *Genome Biol.* 13:R67
122. Taliaferro JM, Lambert NJ, Sudmant PH, Dominguez D, Merkin JJ, et al. 2016. RNA sequence context effects measured in vitro predict in vivo protein binding and regulation. *Mol. Cell* 64:294–306
123. Taliaferro JM, Wang ET, Burge CB. 2014. Genomic analysis of RNA localization. *RNA Biol.* 11:1040–50

124. ten Klooster JP, Leeuwen I, Scheres N, Anthony EC, Hordijk PL. 2007. Rac1-induced cell migration requires membrane recruitment of the nuclear oncogene SET. *EMBO J.* 26:336–45
125. Tian B, Manley JL. 2013. Alternative cleavage and polyadenylation: the long and short of it. *Trends Biochem. Sci.* 38:312–20
126. Treck T, Larson DR, Moldon A, Query CC, Singer RH. 2011. Single-molecule mRNA decay measurements reveal promoter-regulated mRNA stability in yeast. *Cell* 147:1484–97
127. Tsvetanova NG, Klass DM, Salzman J, Brown PO. 2010. Proteome-wide search reveals unexpected RNA-binding proteins in *Saccharomyces cerevisiae*. *PLOS ONE* 5:e12671
128. Tu BP, Kudlicki A, Rowicka M, McKnight SL. 2005. Logic of the yeast metabolic cycle: temporal compartmentalization of cellular processes. *Science* 310:1152–58
129. Ule J, Jensen KB, Ruggiu M, Mele A, Ule A, Darnell RB. 2003. CLIP identifies Nova-regulated RNA networks in the brain. *Science* 302:1212–15
130. Ulitsky I, Shkumatava A, Jan C, Subtelny AO, Koppstein D, et al. 2012. Extensive alternative polyadenylation during zebrafish development. *Genome Res.* 22:2054–66
131. van Meer G, Voelker DR, Feigenson GW. 2008. Membrane lipids: where they are and how they behave. *Nat. Rev. Mol. Cell Biol.* 9:112–24
132. Van Nostrand EL, Pratt GA, Shishkin AA, Gelboin-Burkhart C, Fang MY, et al. 2016. Robust transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). *Nat. Methods* 13:508–14
133. Vourekas A, Alexiou P, Vrettos N, Maragkakis M, Mourelatos Z. 2016. Sequence-dependent but not sequence-specific piRNA adhesion traps mRNAs to the germ plasm. *Nature* 531:390–94
134. Wallace EW, Kear-Scott JL, Pilipenko EV, Schwartz MH, Laskowski PR, et al. 2015. Reversible, specific, active aggregates of endogenous proteins assemble upon heat stress. *Cell* 162:1286–98
135. Walter P, Johnson AE. 1994. Signal sequence recognition and protein targeting to the endoplasmic reticulum membrane. *Annu. Rev. Cell Biol.* 10:87–119
136. Wang ET, Cody NA, Jog S, Biancolella M, Wang TT, et al. 2012. Transcriptome-wide regulation of pre-mRNA splicing and mRNA localization by muscleblind proteins. *Cell* 150:710–24
137. Wang Y, Arribas-Layton M, Chen Y, Lykke-Andersen J, Sen GL. 2015. DDX6 orchestrates mammalian progenitor function through the mRNA degradation and translation pathways. *Mol. Cell* 60:118–30
138. Weber SC, Brangwynne CP. 2012. Getting RNA and protein in phase. *Cell* 149:1188–91
139. Weinberg DE, Shah P, Eichhorn SW, Hussmann JA, Plotkin JB, Bartel DP. 2016. Improved ribosome-footprint and mRNA measurements provide insights into dynamics and regulation of yeast translation. *Cell Rep.* 14:1787–99
140. Wells JN, Bergendahl LT, Marsh JA. 2015. Co-translational assembly of protein complexes. *Biochem. Soc. Trans.* 43:1221–26
141. Whitty A. 2008. Cooperativity and biological complexity. *Nat. Chem. Biol.* 4:435–39
142. Wissink EM, Fogarty EA, Grimson A. 2016. High-throughput discovery of post-transcriptional cis-regulatory elements. *BMC Genom.* 17:177
143. Xie X, Lu J, Kulbokas EJ, Golub TR, Mootha V, et al. 2005. Systematic discovery of regulatory motifs in human promoters and 3' UTRs by comparison of several mammals. *Nature* 434:338–45
144. Yartseva V, Takacs CM, Vejnar CE, Lee MT, Giraldez AJ. 2016. RESA identifies mRNA-regulatory sequences at high resolution. *Nat. Methods* 14:201–7
145. Yeo GW, Coufal NG, Liang TY, Peng GE, Fu XD, Gage FH. 2009. An RNA code for the FOX2 splicing regulator revealed by mapping RNA–protein interactions in stem cells. *Nat. Struct. Mol. Biol.* 16:130–37
146. Zabezhinsky D, Slobodin B, Rapaport D, Gerst JE. 2016. An essential role for COPI in mRNA localization to mitochondria and mitochondrial function. *Cell Rep.* 15:540–49
147. Zaessinger S, Busseau I, Simonelig M. 2006. Oskar allows *nanos* mRNA translation in *Drosophila* embryos by preventing its deadenylation by Smaug/CCR4. *Development* 133:4573–83
148. Zhang H, Elbaum-Garfinkle S, Langdon EM, Taylor N, Ochchipinti P, et al. 2015. RNA controls PolyQ protein phase transitions. *Mol. Cell* 60:220–30
149. Zhao W, Pollack JL, Blagev DP, Zaitlen N, McManus MT, Erle DJ. 2014. Massively parallel functional annotation of 3' untranslated regions. *Nat. Biotechnol.* 32:387–91

150. Zhou HL, Geng C, Luo G, Lou H. 2013. The p97-UBXD8 complex destabilizes mRNA by promoting release of ubiquitinated HuR from mRNP. *Genes Dev.* 27:1046–58
151. Zid BM, O’Shea EK. 2014. Promoter sequences direct cytoplasmic localization and translation of mRNAs during starvation in yeast. *Nature* 514:117–21
152. Zimyanin VL, Belaya K, Pecreaux J, Gilchrist MJ, Clark A, et al. 2008. In vivo imaging of *oskar* mRNA transport reveals the mechanism of posterior localization. *Cell* 134:843–53





## New From Annual Reviews:

### ***Annual Review of Cancer Biology***

cancerbio.annualreviews.org • Volume 1 • March 2017

**ONLINE NOW!**


Co-Editors: **Tyler Jacks**, *Massachusetts Institute of Technology*

**Charles L. Sawyers**, *Memorial Sloan Kettering Cancer Center*

The *Annual Review of Cancer Biology* reviews a range of subjects representing important and emerging areas in the field of cancer research. The *Annual Review of Cancer Biology* includes three broad themes: Cancer Cell Biology, Tumorigenesis and Cancer Progression, and Translational Cancer Science.

#### TABLE OF CONTENTS FOR VOLUME 1:

- *How Tumor Virology Evolved into Cancer Biology and Transformed Oncology*, Harold Varmus 
- *The Role of Autophagy in Cancer*, Naiara Santana-Codina, Joseph D. Mancias, Alec C. Kimmelman
- *Cell Cycle-Targeted Cancer Therapies*, Charles J. Sherr, Jiri Bartek
- *Ubiquitin in Cell-Cycle Regulation and Dysregulation in Cancer*, Natalie A. Borg, Vishva M. Dixit
- *The Two Faces of Reactive Oxygen Species in Cancer*, Colleen R. Reczek, Navdeep S. Chandel
- *Analyzing Tumor Metabolism In Vivo*, Brandon Faubert, Ralph J. DeBerardinis
- *Stress-Induced Mutagenesis: Implications in Cancer and Drug Resistance*, Devon M. Fitzgerald, P.J. Hastings, Susan M. Rosenberg
- *Synthetic Lethality in Cancer Therapeutics*, Roderick L. Beijersbergen, Lodewyk F.A. Wessels, René Bernards
- *Noncoding RNAs in Cancer Development*, Chao-Po Lin, Lin He
- *p53: Multiple Facets of a Rubik's Cube*, Yun Zhang, Guillermina Lozano
- *Resisting Resistance*, Ivana Bozic, Martin A. Nowak
- *Deciphering Genetic Intratumor Heterogeneity and Its Impact on Cancer Evolution*, Rachel Rosenthal, Nicholas McGranahan, Javier Herrero, Charles Swanton
- *Immune-Suppressing Cellular Elements of the Tumor Microenvironment*, Douglas T. Fearon
- *Overcoming On-Target Resistance to Tyrosine Kinase Inhibitors in Lung Cancer*, Ibiayi Dagogo-Jack, Jeffrey A. Engelman, Alice T. Shaw
- *Apoptosis and Cancer*, Anthony Letai
- *Chemical Carcinogenesis Models of Cancer: Back to the Future*, Melissa Q. McCreery, Allan Balmain
- *Extracellular Matrix Remodeling and Stiffening Modulate Tumor Phenotype and Treatment Response*, Jennifer L. Leight, Allison P. Drain, Valerie M. Weaver
- *Aneuploidy in Cancer: Seq-ing Answers to Old Questions*, Kristin A. Knouse, Teresa Davoli, Stephen J. Elledge, Angelika Amon
- *The Role of Chromatin-Associated Proteins in Cancer*, Kristian Helin, Saverio Minucci
- *Targeted Differentiation Therapy with Mutant IDH Inhibitors: Early Experiences and Parallels with Other Differentiation Agents*, Eytan Stein, Katharine Yen
- *Determinants of Organotropic Metastasis*, Heath A. Smith, Yibin Kang
- *Multiple Roles for the MLL/COMPASS Family in the Epigenetic Regulation of Gene Expression and in Cancer*, Joshua J. Meeks, Ali Shilatifard
- *Chimeric Antigen Receptors: A Paradigm Shift in Immunotherapy*, Michel Sadelain



# Contents

Witnessing Genome Evolution: Experimental Reconstruction of Endosymbiotic and Horizontal Gene Transfer <i>Ralph Bock</i> .....	1
The Yeast Genomes in Three Dimensions: Mechanisms and Functions <i>Ken-ichi Noma</i> .....	23
Origin and Evolution of the Universal Genetic Code <i>Eugene V. Koonin and Artem S. Novozhilov</i> .....	45
Regeneration Genetics <i>Chen-Hui Chen and Kenneth D. Poss</i> .....	63
Conditional Degrons for Controlling Protein Expression at the Protein Level <i>Toyoaki Natsume and Masato T. Kanemaki</i> .....	83
Mas-Related G Protein–Coupled Receptors and the Biology of Itch Sensation <i>James Meixiong and Xinzhong Dong</i> .....	103
Mosaicism in Cutaneous Disorders <i>Young H. Lim, Zoe Moscato, and Keith A. Choate</i> .....	123
Transcriptional Regulation in Archaea: From Individual Genes to Global Regulatory Networks <i>Mar Martinez–Pastor, Peter D. Tonner, Cynthia L. Darnell, and Amy K. Schmid</i> ...	143
Regulation by 3'-Untranslated Regions <i>Christine Mayr</i> .....	171
Integration of <i>Agrobacterium</i> T-DNA into the Plant Genome <i>Stanton B. Gelvin</i> .....	195
Genetics and Evolution of Social Behavior in Insects <i>Chelsea A. Weitekamp, Romain Libbrecht, and Laurent Keller</i> .....	219
Human Genetic Determinants of Viral Diseases <i>Adam D. Kenney, James A. Dowdle, Leonia Bozzacco, Temet M. McMichael, Corine St. Gelais, Amanda R. Panfil, Yan Sun, Larry S. Schlesinger, Matthew Z. Anderson, Patrick L. Green, Carolina B. López, Brad R. Rosenberg, Li Wu, and Jacob S. Yount</i> .....	241

Sex Determination in the Mammalian Germline <i>Cassy Spiller, Peter Koopman, and Josephine Bowles</i> .....	265
The Genetics of Plant Metabolism <i>Alisdair R. Fernie and Takayuki Tobge</i> .....	287
Genetic and Structural Analyses of RRNPP Intercellular Peptide Signaling of Gram-Positive Bacteria <i>Matthew B. Neiditch, Glenn C. Capodagli, Gerd Prehna, and Michael J. Federle</i> .....	311
Genetic Networks in Plant Vascular Development <i>Raili Ruonala, Donghwi Ko, and Ykä Helariutta</i> .....	335
Big Lessons from Little Yeast: Budding and Fission Yeast Centrosome Structure, Duplication, and Function <i>Ann M. Cavanaugh and Sue L. Jaspersen</i> .....	361
Noncoding RNAs in Polycomb and Trithorax Regulation: A Quantitative Perspective <i>Leonie Ringrose</i> .....	385
The Relationship Between the Human Genome and Microbiome Comes into View <i>Julia K. Goodrich, Emily R. Davenport, Andrew G. Clark, and Ruth E. Ley</i> .....	413
Combining Traditional Mutagenesis with New High-Throughput Sequencing and Genome Editing to Reveal Hidden Variation in Polyploid Wheat <i>Cristobal Uauy, Brande B.H. Wulff, and Jorge Dubcovsky</i> .....	435
Getting Nervous: An Evolutionary Overhaul for Communication <i>Frederique Varoqueaux and Dirk Fasshauer</i> .....	455
Nucleases Acting at Stalled Forks: How to Reboot the Replication Program with a Few Shortcuts <i>Philippe Pasero and Alessandro Vindigni</i> .....	477
Generation and Evolution of Neural Cell Types and Circuits: Insights from the <i>Drosophila</i> Visual System <i>Michael Perry, Nikos Konstantinides, Filipe Pinto-Teixeira, and Claude Desplan</i> .....	501

## Errata

An online log of corrections to *Annual Review of Genetics* articles may be found at <http://www.annualreviews.org/errata/genet>